

Stroke

American Stroke
AssociationSM

A Division of American
Heart Association



JOURNAL OF THE AMERICAN HEART ASSOCIATION

Efficacy and Safety of Endovascular Cooling After Cardiac Arrest. Cohort Study and Bayesian Approach

Michael Holzer, Marcus Müllner, Fritz Sterz, Oliver Robak, Andreas Kliegel, Heidrun Losert, Gottfried Sodeck, Thomas Uray, Andrea Zeiner and Anton N. Laggner

Stroke published online Jun 8, 2006;

DOI: 10.1161/01.STR.0000227265.52763.16

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214
Copyright © 2006 American Heart Association. All rights reserved. Print ISSN: 0039-2499. Online
ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org>

Subscriptions: Information about subscribing to *Stroke* is online at
<http://stroke.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21202-2436. Phone 410-5280-4050. Fax: 410-528-8550. Email:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/static/html/reprints.html>

Efficacy and Safety of Endovascular Cooling After Cardiac Arrest

Cohort Study and Bayesian Approach

Michael Holzer, MD; Marcus Müllner, MD, MSc; Fritz Sterz, MD; Oliver Robak, MD; Andreas Kliegel, MD; Heidrun Losert, MD; Gottfried Sodeck, MD; Thomas Uray, MD; Andrea Zeiner, MD; Anton N. Laggner, MD

Background and Purpose—Recently 2 randomized trials in comatose survivors of cardiac arrest documented that therapeutic hypothermia improved neurological recovery. The narrow inclusion criteria resulted in an international recommendation to cool only a restricted group of primary cardiac arrest survivors. In this retrospective cohort study we investigated the efficacy and safety of endovascular cooling in unselected survivors of cardiac arrest.

Methods—Consecutive comatose survivors of cardiac arrest, who were either cooled for 24 hours to 33°C with endovascular cooling or treated with standard postresuscitation therapy, were analyzed. Complication data were obtained by retrospective chart review.

Results—Patients in the endovascular cooling group had 2-fold increased odds of survival (67/97 patients versus 466/941 patients; odds ratio 2.28, 95% CI, 1.45 to 3.57; $P < 0.001$). After adjustment for baseline imbalances the odds ratio was 1.96 (95% CI, 1.19 to 3.23; $P = 0.008$). When discounting the observational data in a Bayesian analysis by using a sceptical prior the posterior odds ratio was 1.61 (95% credible interval, 1.06 to 2.44). In the endovascular cooling group, 51/97 patients (53%) survived with favorable neurology as compared with 320/941 (34%) in the control group (odds ratio 2.15, 95% CI, 1.38 to 3.35; $P = 0.0003$; adjusted odds ratio 2.56, 1.57 to 4.17). There was no difference in the rate of complications except for bradycardia.

Conclusion—Endovascular cooling improved survival and short-term neurological recovery compared with standard treatment in comatose adult survivors of cardiac arrest. Temperature control was effective and safe with this device. (*Stroke*. 2006;37:000-000.)

Key Words: brain ■ cardiopulmonary resuscitation ■ heart arrest ■ hypoxia

In industrial countries, per year 36 to 128 per 100 000 inhabitants experience a sudden out-of-hospital cardiac arrest.¹ Unfortunately, survival with favorable neurological recovery is still a rare event after resuscitation.² Recently, 2 randomized trials documented that reducing the body temperature to 32°C to 34°C after successful restoration of spontaneous circulation could substantially improve neurological recovery.^{3,4} Based on these 2 trials international recommendations to cool comatose patients after cardiac arrest attributable to ventricular fibrillation have been formulated.⁵ A further small randomized trial included only patients after asystole or pulseless electrical activity and found a nonsignificant increase in survival and improvement of neurological recovery, indicating that there might be also some beneficial effect in these patients.⁶ Because of narrow inclusion criteria, the generalizability of these trials to an unselected population was limited. The second drawback was the partly insufficient cooling, especially in the European multicenter trial, where

patients were cooled with cold air.⁴ Delays in cooling diminish or even abrogate the beneficial effects of hypothermia in experimental models.^{7,8}

Endovascular cooling is a routine rapid-cooling technique.⁹ A balloon catheter is inserted in a central vein and with a “closed loop” internal cooling circuit, the catheter cools the patient’s blood as it circulates past the catheter.¹⁰ The newer devices have also the advantage of automatic feedback control of patient’s temperature and provide controlled active rewarming.^{9,11–13}

The aim of this cohort study was to investigate the efficacy and safety of endovascular cooling in consecutively admitted survivors of cardiac arrest. We used a Bayesian approach to discount nonrandomized evidence.

Methods

Patients

We evaluated comatose patients after witnessed cardiac arrest and successful restoration of spontaneous circulation who were consec-

Received January 10, 2006; final revision received February 28, 2006; accepted March 27, 2006.

From the Department of Emergency Medicine, Medical University Vienna, Austria.

Correspondence to Prof Dr Fritz Sterz, Universitätsklinik für Notfallmedizin, Medizinische Universität Wien, Währinger Gürtel 18-20/6D, 1090 Wien, Austria. E-mail fritz.sterz@meduniwien.ac.at

© 2006 American Heart Association, Inc.

Stroke is available at <http://www.strokeaha.org>

DOI: 10.1161/01.STR.0000227265.52763.16

utively admitted to an emergency department of a tertiary teaching hospital between August 1991 and November 2004. Data collection and cooling procedure were approved by the local ethics committee. We prospectively recorded all relevant clinical data of these patients in a registry database according to the Utstein guidelines.¹⁴ We excluded patients if they were younger than 18 years, if the cause of cardiac arrest was trauma or severe bleeding, if they had terminal disease, were pregnant, had a pre-existing coagulopathy, had a tympanic membrane temperature below 30°C on admission, received hypothermia with a method other than endovascular cooling, if it lasted longer than 240 minutes from restoration of spontaneous circulation until the initiation of cooling, and if they died within the first 24 hours.

The hypothermia group consisted of patients who were cooled with an endovascular cooling device (Icy catheter and CoolGard 3000; Alsius Corp) with or without the administration of cold fluids in the induction phase of therapeutic hypothermia. All other patients served as controls.

General Management

The resuscitation therapy and postresuscitation care was according to established guidelines.¹⁵ We used a standardized treatment protocol for all patients after cardiac arrest at our institution.¹⁶ Besides analgesia and sedation in all patients, we paralyzed the patients in the endovascular cooling group with rocuronium 0.5 mg/kg initially and used a continuous infusion of 0.5 mg/kg per hour thereafter.

Cooling Procedure

Without delay in patients of the endovascular cooling group, we introduced the cooling device (Icy and CoolGard 3000; Alsius Corp) into the inferior vena cava via a femoral vein and used a foley-catheter with a temperature probe (Ruesch Sensor; Ruesch) for automatic feed-back control of the patient's temperature at 33°C. A number of the patients received 30 mL/kg ice-cold Ringer's lactate during the induction phase of hypothermia. Some of these patients are presented in detail elsewhere.¹⁷ Twenty-four hours after start of cooling we rewarmed the patients with a rate of 0.5°C/hour to 36°C. After the patients had reached normothermia, we kept the temperature below 37°C with usual clinical methods.

Definition of the End Points

As outcome parameter we recorded survival until 30 days after cardiac arrest. We also recorded the neurological performance at 30 days with the cerebral performance category score (CPC).^{14,18} Good neurological recovery was defined as CPC 1 or 2 (the patient is at least alert and has sufficient cerebral function to live independently and work part-time).

For the safety analysis, adverse events for the endovascular cooling group were predefined and recorded prospectively. Controls were selected by frequency matching. Matching criteria were witnessed out-of-hospital ventricular fibrillation cardiac arrest of presumed cardiac cause with duration of cardiac arrest longer than 1 minute. In the control group a chart review was conducted to identify all adverse events. Arrhythmias and bleeding events were recorded between admission and 32 hours. Pneumonia, sepsis, acute renal failure and pancreatitis were recorded within the first 7 days.

Statistical Methods

Continuous variables are given as mean±SD if they were normally distributed or as median and the range between the 25th and 75th quartile if they were not. Nominal data are given as counts and percent of total number. We performed univariate comparisons between groups with a χ^2 test or Fisher exact test when indicated. For safety analyses we used contingency tables and Fisher exact tests. We did not correct for multiple testing.

Logistic Regression for Observational Data

We used a logistic regression model providing an approximate normal likelihood for the odds ratio while adjusting for baseline imbalances. We entered the following baseline variables in the

model: age, history of diabetes, New York Heart Association score before arrest, location out-of-hospital arrest, presumed cardiac cause, ventricular fibrillation or ventricular tachycardia as primary rhythm, basic life-support, no-flow, low-flow. We used standard procedures to check proportionality of odds and model-fit.

Bayesian Analysis

In a retrospective Bayesian analysis the observed data (the likelihood) modify a defined prestudy probability (the prior distribution) to obtain an updated probability (the posterior distribution) of the possible values for the variable of interest. This represents a powerful mechanism for incorporating information from previous studies and for controlling confounding.¹⁹ Additionally, randomization is incidental, and it is possible to assess any magnitude of therapeutic response with this method.²⁰

From the posterior probability distributions, means and 95% credible intervals were calculated. A 95% credible interval includes the true value of the variable of interest with 95% probability. We discounted the cohort study data by assuming that the observed effect was actually 0 and probability of exceeding the observed effect was 5%.

The formulas used to calculate the posterior mean and variance of the ln(OR) were²¹:

$$M_1 = M_0 \frac{\frac{1}{V_0}}{\frac{1}{V_0} + \frac{1}{V}} + M \frac{\frac{1}{V}}{\frac{1}{V_0} + \frac{1}{V}} \quad V_1 = \frac{1}{\frac{1}{V_0} + \frac{1}{V}}$$

where M_1 indicates posterior mean; M , data mean; M_0 , prior mean, V_1 , posterior variance; V_0 , prior variance; V , data variance.

We calculated also the magnitude of therapeutic response for a threshold of benefit of 10%, 20% and 30% using the skeptical prior to discount the observational data. This magnitude of therapeutic response gives the respective probability that the risk reduction exceeds a putative "threshold of benefit" given the observational data.²⁰

For all statistical calculations we used Stata, release 8.2 (StataCorp) and MS Excel 2002 (Microsoft). A $P < 0.05$ was considered statistically significant.

Results

Patients

Between August 1991 and November 2004, 1038 patients fulfilled the inclusion and exclusion criteria and were included in the analysis. Of the included patients, 97 were treated with endovascular cooling and 941 served as controls (supplemental Figure I, available online at <http://stroke.ahajournals.org>). The baseline demographic information of all patients is given in Table 1.

Cooling Procedure

Continuous temperature and cooling data were available only for the endovascular cooling group. Admission tympanic temperature was 35.4±1.1°C at 40 (interquartile range [IQR] 29 to 58) minutes after cardiac arrest. Of the 97 patients who were cooled with endovascular cooling, 41 received additional cold Ringer's lactate during induction of hypothermia. Because this might influence the calculated cooling rate of endovascular cooling, we restricted the following analysis to patients who received endovascular cooling exclusively.

The bladder temperature at start of cooling 95 (IQR 67 to 156) minutes after restoration of spontaneous circulation was 35.3±1.0°C. It lasted 253 (IQR 170 to 345) minutes from

TABLE 1. Demographic and Resuscitation Data of the Patients in the Endovascular Cooling and Control Group

	Endovascular Cooling (n=97)	Control (n=941)	P
Female, n (%)	30 (31)	329 (35)	0.426
Age, y (IQR)	52 (42–62)	60 (50–70)	<0.001
Diabetes mellitus, n (%)	9 (9)	179 (19)	0.018
New York Heart Association Class	1 (1–1)	1 (1–2)	<0.001
Out-of-hospital cardiac arrest, n (%)	89 (92)	608 (65)	<0.001
Cardiac cause of cardiac arrest, n (%)	82 (85)	600 (64)	<0.001
Ventricular fibrillation, n (%)	69 (71)	435 (46)	<0.001
BLS, n (%)	41 (42)	240 (26)	<0.001
Time to initiation of BLS (no flow), min (IQR)	5 (1–9)	0 (0–5)	<0.001
Time from initiation of BLS to ROSC (low flow), min (IQR)	15 (9–25)	13 (5–22)	0.0734
Rearrest within 24 h, n (%)	12 (12)	153 (16)	0.319
Survival at 1 mo, n (%)	67 (69)	466 (50)	<0.001
Survival and Good Neurology (CPC 1 or 2) at 1 mo, n (%)	51 (53)	320 (34)	<0.001

IQR indicates range between the 25th and 75th quartile; No flow, time interval from cardiac arrest to first resuscitation attempts; ROSC, restoration of spontaneous circulation; BLS, basic life-support.

restoration of spontaneous circulation to reach the target temperature of 33°C. The cooling rate was 1.2 (IQR 0.7 to 1.5)°C/hour with endovascular cooling. The patients were cooled for 24.3 (IQR 22.0 to 25.5) hours. Thereafter, they were rewarmed within 388 (IQR 313 to 579) minutes to a temperature of 36°C (Figure 1).

In 14 of the 97 patients, cooling was terminated after median 11.7 (min 5.7; max 19.1) hours after start of cooling attributable to hemodynamic instability (n=2), bleeding at the catheter insertion site (n=2), transfer to the operation theater (n=2), obvious signs of brain death (n=1) and various

other causes (n=7) including multiple organ failure and subarachnoid hemorrhage.

Outcome

Regardless of treatment, 533 of 1038 patients (51%) survived for 30 days and 371 of 1038 patients (36%) survived for 30 days and had favorable neurological recovery (CPC 1 or 2).

Survival to 30 Days

In the univariate analysis patients in the endovascular cooling group had 2-fold increased odds of survival (67 of 97 patients versus 466 of 941 patients; odds ratio 2.28, 95% CI, 1.45 to 3.57; *P*<0.001). After adjustment for baseline imbalances in a multivariate model the odds ratio was only slightly attenuated (odds ratio 1.96, 95% CI, 1.19 to 3.23; *P*=0.008). The model had an acceptable fit (Hosmer-Lemeshow $\chi^2=7.39$, *df*=8; *P*=0.495). When discounting the observational data in a Bayesian analysis by using a skeptical prior the posterior odds ratio was 1.61 (95% credible interval, 1.06 to 2.44; Figure 2). The calculated magnitude of therapeutic response for a threshold of benefit of 10%, 20% and 30% using the skeptical prior to discount the observational data is shown in Table 2.

Survival to 30 Days or Until Discharge and Good Neurological Recovery

In the univariate analysis 51/97 patients (53%) in the endovascular cooling group survived with favorable neurology as compared with 320/941 (34%) in the control group (odds ratio 2.15, 95% CI, 1.38 to 3.35; *P*=0.0003). After adjustment for baseline imbalances in a multivariate model, the odds ratio was further increased (odds ratio 2.56, 95% CI, 1.57 to 4.17; *P*<0.001). The model had an acceptable fit (Hosmer-Lemeshow $\chi^2=8.22$, *df*=8; *P*=0.413).

Safety of Endovascular Cooling

There were no statistically significant differences in adverse events between the frequency-matched groups except for the occurrence of transient bradycardia (Table 3). In most cases the rate of adverse events was comparable. Transient increases in renal (n=11) and pancreatic enzymes (n=4) attended the period of cooling and resolved with rewarming.




Figure 1. Bladder temperature course. Median, 25th and 75th quartile of bladder temperature after return of spontaneous circulation in patients, who were exclusively cooled with the endovascular cooling device (n=56). Target temperature, 33°C; cooling duration, 24 hours.




Figure 2. Discounted odds ratio of observational data. Odds ratio (log-scale) for observational data by applying a skeptical prior (assuming that the observed effect is actually 0 and probability of exceeding the observed effect was 5%). Dotted line, skeptical prior; dashed line, observational data; continuous line, posterior odds ratio (discounted data).

Dialysis or other clinical interventions were not required. Attributable to bleeding at the insertion site of the endovascular cooling catheter, cooling was terminated in 2 cases (5 and 14 hours after start of cooling). One of the 2 cases received thrombolysis, the other underwent coronary angioplasty. However, the total number of bleeding events was not different between the groups.

Discussion

Treatment of unselected patients after resuscitation from cardiac arrest with the Icy-CoolGard3000 cooling device significantly reduced mortality and improved favorable neurological recovery at 30 days or discharge as compared with controls from our resuscitation database. Therapeutic hypothermia with endovascular cooling was not associated with a higher rate of adverse events.

Based on 2 randomized controlled trials,^{3,4} mild therapeutic hypothermia both reduced neurological damage and improved survival in primary survivors of cardiac arrest. The discounted odds ratio obtained in the Bayesian analysis of 1.61 (95% credible interval, 1.06 to 2.44) found in our study compared well to the calculated odds ratios of 1.77 (95% CI, 1.07 to 2.94)⁴ and 2.00 (95% CI, 0.71 to 5.69)³ in the 2 large randomized trials. In contrast to these studies, consecutive patients with various causes of arrest and with other primary rhythms than ventricular fibrillation or ventricular tachycardia were included in our trial. This makes the results obtained by our study more generalizable and suggests that hypothermia will be the therapy of choice for all

comatose patients who are successfully resuscitated from cardiac arrest.

The cooling rate of 1.2°C/hour in our study compared well with reported cooling rates of endovascular cooling in stroke,¹² subarachnoid hemorrhage,¹³ myocardial infarction^{11,22} and cardiac arrest,⁹ which was between 0.8°C/hour and 4.8°C/hour. This rate was much faster than external cooling with a water-filled cooling blanket²³ or with cold air⁴ where 14% of the patients never reached target temperature, 70% required additional ice packs and the mean time to target temperature was 480 minutes.

Recently, a rapid method of induction of hypothermia was studied by Bernard et al.²⁴ The infusion of 30 mL/kg of ice-cold (4°C) intravenous lactated Ringer's solution over 30 minutes achieved a rapid decrease in median core temperature from 35.5°C to 33.8°C (cooling rate 3.4°C/hour) in 22 comatose survivors of out-of-hospital cardiac arrest. The combination of these 2 techniques would induce hypothermia

TABLE 2. Posterior Probability of Benefit of Empirical Data Discounted With a Sceptical Prior

Threshold of Benefit, %	Posterior Probability of Benefit of Empirical Data Discounted With a Sceptical Prior
10	0.96
20	0.92
30	0.84

TABLE 3. Complications During and After Endovascular Cooling Compared to Frequency-Matched Controls

Adverse Event	Endovascular		P
	Cooling (n=62)	Control (n=104)	
Within the first 32 h			
Atrial fibrillation, n (%)	2 (3)	2 (3)	0.987
Ventricular tachycardia, n (%)	14 (23)	9 (14)	0.231
Ventricular fibrillation, n (%)	6 (10)	6 (10)	0.977
Narrow complex tachycardia, n (%)	0	3 (5)	0.082
Bradycardia, n (%)	9 (15)	2 (3)	0.025
Any Bleeding, n (%)	16 (26)	27 (26)	0.982
Within the first 7 d			
Pneumonia, n (%)	17 (27)	20 (19)	0.233
Elevation of pancreatic enzymes, n (%)	1 (2)	0	0.194
Sepsis, n (%)	0	0	...
Acute renal failure, n (%)	4 (6)	4 (4)	0.448

rapidly and then allows a convenient control of the temperature over the selected time by endovascular cooling. A further advantage of endovascular cooling is the possibility of an individual rewarming speed which could be adapted to various etiologies of ischemic damage.

There was no statistically significant difference in adverse events between the endovascular cooling group and the control group except for bradycardia (Table 3). The elevated number of minor bleedings at the cooling catheter insertion site could be explained by the use of anticoagulation in patients who had a myocardial infarction and underwent thrombolysis or angioplasty. Pneumonia and sepsis might be a further problem as suggested by the European multicenter trial of therapeutic hypothermia after cardiac arrest.⁴ These complications did not occur more frequently in the endovascular cooling group in our study. If this was caused by the more precise temperature management attributable to feedback, control of bladder temperature is not known, but there was virtually no overshoot during induction of hypothermia (Figure 1).

There are several limitations to mention. This trial was not blinded for the assessment of neurological outcome, which might have led to information bias. This is not an issue, however, for the end point of survival. Furthermore, only few patients with unfavorable neurological recovery survive hospital discharge,⁴ and “Do Not Resuscitate” orders might also have an impact on outcome. Because this was a retrospective cohort study, selection bias might have occurred. Although there has been a department policy of cooling all patients after cardiac arrest, who fulfilled the inclusion and exclusion criteria, the final decision to cool a patient was left to the discretion of the attending physician. Therefore, it is possible that a higher proportion of patients with expected bad outcome have not been cooled. There were imbalances in prognostic baseline variables. We tried to adjust for these imbalances by using standard multivariate analyses. We like to point out that the outcome was only moderately confounded by the baseline imbalances. A further problem might be that the exact course of the resuscitation can only be estimated.²⁵ We tried to reduce this problem by thoroughly interviewing all involved individuals immediately after resuscitation and entering all data in a central database. All of these limitations could have led to an overestimation of effect size. This was accounted for by further discounting the result by combining it with a skeptical prior in a Bayesian analysis. Interestingly however, the observed effect on neurological outcome and survival of endovascular cooling had the same magnitude as with other cooling methods used in randomized controlled trials.^{3,4}

Endovascular cooling via the Icy-CoolGard cooling device improved survival and short-term neurological recovery compared with standard treatment in adult survivors of cardiac arrest. Temperature control was effective and safe with this device.

Source of Funding

This work was supported by a grant (grant number: 10343) of the “Jubilaeumsfond” of the Oesterreichische Nationalbank, Vienna, Austria.

Disclosures

Alsius Corp was not involved in the design of the study, data management, data analysis, or manuscript preparation or authorship. Alsius Corp was allowed to review the manuscript. However, any decisions regarding manuscript revision were made by the authors. Alsius Corp provided travel grants for scientific meetings for M.H., A.K., and F.S. T.U. and G.S. were employed for 6 months at the Department of Emergency Medicine, Medical University Vienna with support from a grant from Alsius Corp. H.L. received a laptop from Alsius Corp. O.R. received payment for data extraction from patient’s charts on a ‘by the hour’ basis from Alsius Corp.

M.M., A.Z. and A.N.L. have no conflicts of interest to declare.

References

1. Vreede-Swagemakers JJ, Gorgels AP, Dubois-Arbouw WI, van Ree JW, Daemen MJ, Houben LG, Wellens HJ. Out-of-hospital cardiac arrest in the 1990’s: a population-based study in the Maastricht area on incidence, characteristics and survival. *J Am Coll Cardiol.* 1997;30:1500–1505.
2. Herlitz J, Bahr J, Fischer M, Kuisma M, Lexow K, Thorgeirsson G. Resuscitation in Europe: a tale of five European regions. *Resuscitation.* 1999;41:121–131.
3. Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, Smith K. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med.* 2002;346:557–563.
4. The Hypothermia After Cardiac Arrest (HACA) study group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med.* 2002;346:549–556.
5. Nolan JP, Morley PT, Hoek TL, Hickey RW. Therapeutic hypothermia after cardiac arrest. An advisory statement by the Advancement Life support Task Force of the International Liaison Committee on Resuscitation. *Resuscitation.* 2003;57:231–235.
6. Hachimi-Idrissi S, Corne L, Ebinger G, Michotte Y, Huyghens L. Mild hypothermia induced by a helmet device: a clinical feasibility study. *Resuscitation.* 2001;51:275–281.
7. Kuboyama K, Safar P, Radovsky A, Tisherman SA, Stezoski SW, Alexander H. Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: a prospective, randomized study. *Crit Care Med.* 1993;21:1348–1358.
8. Colbourne F, Corbett D. Delayed postischemic hypothermia: a six month survival study using behavioral and histological assessments of neuroprotection. *J Neurosci.* 1995;15:7250–7260.
9. Al Senani FM, Graffagnino C, Grotta JC, Saiki R, Wood D, Chung W, Palmer G, Collins KA. A prospective, multicenter pilot study to evaluate the feasibility and safety of using the CoolGard System and Icy catheter following cardiac arrest. *Resuscitation.* 2004;62:143–150.
10. Behmann FW, Bontke E. Intravasale Kühlung; eine Methode zur Erzielung steuerbarer Hypothermie. *Pflugers Arch.* 1956;263:145–165.
11. Dixon SR, Whitbourn RJ, Dae MW, Grube E, Sherman W, Schaer GL, Jenkins JS, Baim DS, Gibbons RJ, Kuntz RE, Popma JJ, Nguyen TT, O’Neill WW. Induction of mild systemic hypothermia with endovascular cooling during primary percutaneous coronary intervention for acute myocardial infarction. *J Am Coll Cardiol.* 2002;40:1928–1934.
12. Georgiadis D, Schwarz S, Kollmar R, Schwab S. Endovascular cooling for moderate hypothermia in patients with acute stroke: first results of a novel approach. *Stroke.* 2001;32:2550–2553.
13. Keller E, Imhof HG, Gasser S, Terzic A, Yonekawa Y. Endovascular cooling with heat exchange catheters: a new method to induce and maintain hypothermia. *Intensive Care Med.* 2003;29:939–943.
14. Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, Bossaert L, Deloos HH, Dick WF, Eisenberg MS. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation.* 1991;84:960–975.
15. European Resuscitation Council. International Guidelines 2000 for CPR and ECC. A consensus on science. *Resuscitation.* 2000;46:1–447.

16. Kliegel A, Havel C, Sterz F. Die Behandlung des Patienten mit Herz-Kreislaufstillstand und nach Reanimation im Krankenhaus. *Intensivmedizin und Notfallmedizin*. 2002;39:13–25.
17. Kliegel A, Losert H, Sterz F, Kliegel M, Holzer M, Uray T, Domanovits H. Cold simple intravenous infusions preceding special endovascular cooling for faster induction of mild hypothermia after cardiac arrest—a feasibility study. *Resuscitation*. 2005;64:347–351.
18. Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet*. 1975;1:480–484.
19. Dunson DB. Commentary: practical advantages of Bayesian analysis of epidemiologic data. *Am J Epidemiol*. 2001;153:1222–1226.
20. Diamond GA, Kaul S. Prior convictions: Bayesian approaches to the analysis and interpretation of clinical megatrials. *J Am Coll Cardiol*. 2004;43:1929–1939.
21. Lee PM. Bayesian inference for the normal distribution. In: *Bayesian Statistics: An Introduction*. 3rd edition. New York, NY: Oxford University Press; 2004: 31.
22. Kandzari DE, Chu A, Brodie BR, Stuckey TA, Hermiller JB, Vetrovec GW, Hannan KL, Krucoff MW, Christenson RH, Gibbons RJ, Sigmon KN, Garg J, Hasselblad V, Collins K, Harrington RA, Berger PB, Chronos NA, Hochman JS, Califf RM. Feasibility of endovascular cooling as an adjunct to primary percutaneous coronary intervention (results of the LOWTEMP pilot study). *Am J Cardiol*. 2004;93:636–639.
23. Felberg RA, Krieger DW, Chuang R, Persse DE, Burgin WS, Hickenbottom SL, Morgenstern LB, Rosales O, Grotta JC. Hypothermia after cardiac arrest: feasibility and safety of an external cooling protocol. *Circulation*. 2001;104:1799–1804.
24. Bernard S, Buist M, Monteiro O, Smith K. Induced hypothermia using large volume, ice-cold intravenous fluid in comatose survivors of out-of-hospital cardiac arrest: a preliminary report. *Resuscitation*. 2003;56:9–13.
25. Kaye W, Mancini ME, Truitt TL. When minutes count—the fallacy of accurate time documentation during in-hospital resuscitation. *Resuscitation*. 2005;65:285–290.



Stroke

JOURNAL OF THE AMERICAN HEART ASSOCIATION

FIRST PROOF ONLY




Figure 1. Flow of participants. All patients in the database from August 1991 to November 2004 were screened. For outcome evaluation all patients who were cooled with endovascular cooling during this period were evaluated. For evaluation of cooling rate we restricted the analysis to patients who received endovascular cooling exclusively.

American Stroke Association
A Division of American Heart Association

Stroke

JOURNAL OF THE AMERICAN HEART ASSOCIATION

FIRST PROOF ONLY